

A

## ANSWER KEY

1) B	2) A	3) C	4) B	5) C
6) D	7) A	8) A	9) D	10) A
11) B	12) D	13) C	14) C	15) B
16) B	17) D	18) C	19) B	20) B
21) B	22) B	23) D	24) C	25) B
26) D	27) A	28) B	29) A	30) A
31) A	32) B	33) A	34) B	35) A
36) C	37) A	38) B	39) B	40) C
41) B	42) C	43) B	44) C	45) B
46) C	47) D	48) D	49) A	50) A
51) B	52) C	53) C	54) C	55) D
56) C	57) B	58) C	59) B	60) B
61) C	62) B	63) B	64) D	65) D
66) A	67) C	68) D	69) D	70) C
71) D	72) B	73) A	74) C	75) C
76) A	77) B	78) A	79) B	80) D
81) B	82) C	83) A	84) C	85) D
86) C	87) A	88) A	89) C	90) D
91) A	92) A	93) D	94) A	95) C
96) A	97) C	98) B	99) D	100) C

HINTS & SOLUTION

1. (b) Median =  $\frac{5\text{th Term} + 6\text{th Term}}{2}$   
 $\Rightarrow 61.5 = \frac{x + x + 3}{2}$   
 $\Rightarrow 123 = 2x + 3$   
 $\Rightarrow x = 60$

2. (a)

X	f	fx
0	4	0
1	f	f
2	9	18
3	g	3g
4	4	16

$\sum fx = f + 3g + 34$ ,  $N = 25$ ,  $\bar{X} = 2$

$f + g + 17 = 25 \Rightarrow f + g = 8$

$\bar{X} = \frac{\sum fx}{N}$

$2 = \frac{f + 3g + 34}{25}$

$50 = (f + g) + 2g + 34$

$50 = 8 + 2g + 34$

$g = 4, f = 4$

3. (c) Because there is a gap between two adjacent bars, so both the districts can be represented by bar chart.

4. (a) Let the frequency of class interval 21-30 be f

$9 + 22 + f + 20 + 12 + 8 = 100$

$\Rightarrow 71 + f = 100 \Rightarrow f = 29$

Highest frequency = 29

$\therefore$  Modal class = 20.5 – 30.5


5. (a)

I.  $HM = \frac{2 \times 8 \times 12}{8 + 12} = \frac{48}{5} = 9.6$

II.  $HM = \frac{2 \times 9 \times 11}{9 + 11} = \frac{198}{20} = 9.9$

III.  $HM = \frac{2 \times 6 \times 24}{6 + 24} = \frac{288}{30} = 9.6$

6. (d) Join OB.  $OA = OB = OC$



Then,  $\angle OAB = \angle OBA = 20^\circ$


$\angle OCB = \angle OBC = 30^\circ$

$\angle ABC = 50^\circ$

We know that  $\angle ABC = \frac{1}{2} \angle AOC$

$\angle AOC = 2 \angle ABC = 2 \times 50^\circ = 100$

7. (a)



$r_1 + r_2 = 4, r_2 + r_3 = 6, r_3 + r_1 = 8$

$$\begin{aligned} &(r_1 + r_2) + (r_2 + r_3) + (r_3 + r_1) \\ &\Rightarrow 4 + 6 + 8 \\ &2(r_1 + r_2 + r_3) = 18 \\ &r_1 + r_2 + r_3 = \frac{18}{2} = 9 \end{aligned}$$

8. (a) We know that, the triangle of same segment of a circle makes an equal angles.

$$\angle XBY = \angle XAY = 45^\circ$$

$$\text{In } \triangle BXY, \angle BXY + \angle XBY + \angle BYX = 180^\circ$$

$$\Rightarrow 50^\circ + 45^\circ + \angle BYX = 180^\circ$$

$$\Rightarrow \angle BYX = 180^\circ - 95^\circ = 85^\circ$$

9. (a)

$$\angle BOD = 180^\circ - 106^\circ = 74^\circ$$

Since,  $\angle BOD$  is an angle made by arc BD on centre.

Here,  $\angle BCD$  is an angle made by arc BD on circumference

$$\angle BCD = \frac{1}{2} \angle BOD$$

$$= \frac{1}{2} \times 74^\circ = 37^\circ$$

10. (a) Only one circle passing through all the vertices of a given triangle.

11. (b) Area of Parallelogram =  $6 \times$  Area of  $\triangle NPR$

$$NR \times PL = 6 \times \frac{1}{2} \times NR \times PR$$

$$PL = 3 PR$$


$$PR + RL = 3PR$$

$$RL = 2PR = 2 \times 6 = 12$$


12. (d) If two parallel lines are cut by two distinct transversals, the quadrilateral formed by the four lines is always a

‘Trapezium’.

**Case I** If two distinct transversals (are not parallel), then always  $\rightarrow$  (Trapezium)




**Case II** If two distinct transversals are parallel, then always (Trapezium + Parallelogram)



13. (c) Join AC

In  $\triangle ACD$ ,  $EG \parallel DC$ , E & G are mid points of AD and AC respectively



$$EG = \frac{1}{2} DC \Rightarrow EG = \frac{3}{2}$$

$$GF = \frac{1}{2} AB \Rightarrow GF = 1$$

$$EF = EG + GF = 1 + \frac{3}{2} = \frac{5}{2}$$

$$\text{Ratio} = \frac{\text{Area of } ABFE}{\text{Area of } EFCD}$$

$$= \frac{\frac{1}{2} \left( 2 + \frac{5}{2} \right) \times h}{\frac{1}{2} \left( 3 + \frac{5}{2} \right) \times h} = \frac{9}{11}$$

14. (c) Given,  $BC \parallel EF \parallel AD$

$x^\circ = z^\circ = 50^\circ$  (corresponding interior angle)

$$\theta + z^\circ = 180^\circ \text{ (Linear pair)}$$

$$\theta = 180^\circ - 50^\circ = 130^\circ$$


In Quadrilateral

$$AQFD, x^\circ + y^\circ + 120^\circ + \theta = 360^\circ$$

$$50^\circ + y^\circ + 120^\circ + 130^\circ = 360^\circ$$

$$y = 360^\circ - 300^\circ = 60^\circ$$

15. (b) Since, line segment  $AP, CQ$  bisect the  $\angle A, \angle C$  respectively. Then,  $AP \parallel CQ$



In  $\Delta APQ$  &  $\Delta CPQ$ ,  
 $AP \parallel CQ$

$$\angle APQ = \angle PQC$$

$PQ$  is common


$$PC \parallel AQ$$

Hence,  $\angle CPQ = \angle PQA$  (Alternate angles)

$\Delta APQ \sim \Delta CQP$  (by ASA)

$$\therefore \Delta APQ \sim \Delta PCQ$$

16. (d)




$$AB = 3AP$$

In  $\Delta ABC$  and  $\Delta APQ$ ,  $PQ \parallel BC$  and  $\angle A$  is common. Therefore,  $\Delta ABC \sim \Delta APQ$

$$\Rightarrow \frac{\text{Area of } \Delta APQ}{\text{Area of } \Delta ABC} = \frac{AP^2}{AB^2} = \left(\frac{1}{3}\right)^2 = \frac{1}{9}$$

17. (c) G is a centroid of triangle



In  $\Delta ABC$ ,  $AD$  is median of  $\Delta ABC$


$AG : GD = 2 : 1$  (By property of triangle)

$$\Rightarrow 8 : GD = 2 : 1$$

$$\Rightarrow GD = 4 \text{ cm}$$

Therefore,  $AD = 8 + 4 = 12$

18. (b)



$$AC^2 = AB^2 + BC^2$$


$$\Rightarrow AC^2 = 64 + 64$$

$$\Rightarrow AC^2 = 8\sqrt{2}$$

$ABC$  is isosceles right angle triangle

$$AP = PC = PB = \frac{AC}{2} = 4\sqrt{2}$$

19. (b)



In  $\Delta ABC$ ,  $\angle ABC = \angle CAB = 38^\circ$

$$\angle ACB = 180^\circ - (\angle ABC + \angle CAB)$$

$$\Rightarrow 180^\circ - (38^\circ + 38^\circ) = 104^\circ$$

In  $\Delta ACD$ ,  $\angle ACD = 180^\circ - 104^\circ = 76^\circ$

$$CD = AD, \therefore \angle ACD = \angle CAD = 76^\circ$$


$$\angle ADC = 180^\circ - (\angle ACD + \angle CAD)$$

$$\Rightarrow 180^\circ - (76^\circ + 76^\circ) = 28^\circ$$

20. (b) Let the height of triangle be  $x$ , then

$$BC = 7x$$

$$\text{Area of } \Delta ABC = \frac{1}{2} \times 7x \times x = \frac{7}{2}x^2$$



Cost of painting the wall at ₹350 per 100 sq m = ₹1225

Cost of painting 100 m<sup>2</sup> = ₹350

Cost of painting 1 m<sup>2</sup> = ₹3.5

$$\text{Cost of painting } \frac{7}{2}x^2 = \frac{350}{100} \times \frac{7}{2}x^2$$

$$\Rightarrow \frac{350}{100} \times \frac{7}{2}x^2 = 1225$$

$$\Rightarrow x^2 = \frac{1225 \times 100 \times 2}{350 \times 7} = \frac{35 \times 10 \times 2}{7}$$

$$\Rightarrow 5 \times 10 \times 2 = 100 = 10 m$$

$$\text{Base} = BC = 7x = 70$$

21. (b)  $\angle PTB = \angle ATU = 55^\circ$

(Vertically opposite angles)

Similarly,  $\angle DVS = \angle BTV = 45^\circ$

(Corresponding angles)

$$\angle RTP = 180^\circ - (45^\circ + 55^\circ) = 80^\circ$$


$$\angle PTB = \angle TUV = 55^\circ$$

(Corresponding angles)


$$\angle CUQ = \angle TUV = 55^\circ$$

(Vertically opposite angles)

$$\text{Sum of angles} = \angle RTP + \angle CUQ = 135^\circ$$



22. (b) Given  $AC^2 = AB \times CB$



$$\Rightarrow x^2 = 2 \times (2 - x)$$

$$\Rightarrow x^2 = 4 - 2x$$

$$x^2 + 2x - 4 = 0$$

$$x = \frac{-2 \pm \sqrt{4 + 16}}{2 \times 1}$$

$$x = -1 \pm \sqrt{5}$$


$$BC = 2 - (-1 \pm \sqrt{5})$$

$$= 3 - \sqrt{5}$$


23. (d) We know that, when two lines intersect each other it makes 4 angles.

The total number of pairs = 3

Total number of angles =  $3 \times 4 = 12$



24. (c) Let us draw a line parallel to AB which is EF.



$$\angle EOP = \angle OPB$$

$$\Rightarrow \angle EOP = b$$

$$\angle EON = \angle ONM$$


$$\Rightarrow \angle EON = c$$

$$\angle PON = b + c$$

$$\angle PON + a = 2\pi$$

$$\Rightarrow a = 2\pi - (b + c)$$

25. (b) Here  $r_1 = 3\text{cm}$ ,  $r_2 = 6\text{cm}$ ,  $h = 9\text{cm}$ ,



Total surface area of the frustum

$$\pi [(R + r)l^2 + r^2 + R^2]$$

Where,  $l = \sqrt{h^2 + (R - r)^2}$

$$\Rightarrow \pi [(6 + 3)\sqrt{81 + 9 + 9 + 36}]$$

$$\Rightarrow \pi [9\sqrt{90 + 45}] = 9\pi [3\sqrt{10 + 5}]$$

26. (d) By using properties at similar triangle in  $\Delta MPA, \Delta MOC$

$$\frac{MP}{MO} = \frac{PA}{OC}$$

$$\Rightarrow \frac{x}{9 + x} = \frac{3}{6} \Rightarrow x = 9$$

Height of the cone =  $MO = x + 9$   
 $= (9 + 9) = 18\text{cm}$

27. (a) In  $\Delta ABC$

$$AC^2 = AB^2 + BC^2$$

$$\Rightarrow AC^2 = 9^2 + (6 - 3)^2 = 90$$

$$\Rightarrow AC = 3\sqrt{10}$$

28. (b) Given,  $l = 30\text{ cm}, b = 24\text{cm}, h = 18\text{cm}$

Maximum length of the rod can be placed in the cuboid is equal to diagonal of box


$$\sqrt{30^2 + 24^2 + 18^2} = \sqrt{900 + 576 + 324}$$

$$\Rightarrow \sqrt{1800} = 30\sqrt{2}$$

29. (a) Increase in the height of water level

$$\frac{0.75}{2.5 \times 1.5} = 0.2\text{m or } 20\text{ cm}$$

30. (a) Here width of sheet is 20 cm, which is the maximum diameter of the circular sheet.



Remaining area of sheet = Area of rectangle sheet – Area of circular sheet  
 $= 25 \times 20 - \pi(10)^2 = 500 - 314 = 186$

31. (a) Circumference of circle =  $2\pi \times 42$

$$2 \times \frac{22}{7} \times 42 = 264$$

Perimeter of square =  $4x$

$$\Rightarrow 4x = 264. \quad x = 66$$

32. (b) The above figure is symmetrical about BD

Area of shaded part

$$= 2 \times \text{Area of BEDB}$$

$$= 2 \times \text{Area of BCDEB} - \text{Area of } \Delta BCD$$

$$\Rightarrow 2 \left( \frac{\pi r^2}{4} - \frac{1}{2} \times BC \times CD \right)$$

$$\Rightarrow 2 \left( \frac{22}{7} \times \frac{7^2}{4} - \frac{1}{2} \times 7 \times 7 \right) = 28\text{ cm}$$

33. (a) Let sides of rectangle be  $l$  and  $b$

$$\text{Then } 2(l + b) = 18$$

$$\Rightarrow l + b = 9$$

$$\text{Area of rectangle} = l \times b$$

$$\text{For maximum area, } l = b$$


$$2b = 9, b = 4.5$$

$$\text{Maximum area} = 4.5^2 = 20.25$$

34. (b) Given that  $\beta = 30^\circ$

In  $\Delta AED, \tan \beta = \tan 30^\circ$

$$\tan 30^\circ = \frac{AE}{DE} = \frac{1}{\sqrt{3}}$$



$$DE = \sqrt{3} AE = \sqrt{3} \left( \frac{3 + \sqrt{3}}{2} \right) h$$

$$\Rightarrow BC = DE = \frac{3}{2} (1 + \sqrt{3}) h$$

Now, in  $\Delta CB$

$$\tan \alpha = \frac{AC}{BC}$$

$$\Rightarrow BC \tan \alpha = AE - CE = AE - BD$$

$$\Rightarrow BC \tan \alpha = \left( \frac{3 + \sqrt{3}}{2} \right) h - h = h \left( \frac{3 + \sqrt{3} - 2}{2} \right)$$

$$\Rightarrow \frac{3}{2} (1 + \sqrt{3}) h \tan \alpha = \left( \frac{1 + \sqrt{3}}{2} \right) h$$

$$\tan \alpha = \frac{1}{3}$$

35. (a) Given that  $\alpha = 30^\circ$

In  $\Delta ACB$ ,  $\tan \alpha = \tan 30^\circ$

$$\tan 30^\circ = \frac{AC}{BC} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow BC = \sqrt{3} AC = \sqrt{3} (AE - CE)$$

$$\Rightarrow \sqrt{3} (AE - BD)$$

$$\Rightarrow \sqrt{3} \left( \frac{3 + \sqrt{3}}{2} - 1 \right) h = \frac{\sqrt{3}}{2} (1 + \sqrt{3})$$

Now, in  $\Delta AED$

$$\tan \beta = \frac{AE}{DE} = \frac{AE}{BC}$$

$$\Rightarrow \frac{\left( \frac{3 + \sqrt{3}}{2} \right) h}{\frac{\sqrt{3}}{2} (1 + \sqrt{3}) h} = \frac{\sqrt{3} (1 + \sqrt{3}) h}{2}$$

$$\tan \beta = 1$$

36. (c) Given,  $\alpha = 30^\circ, h = 30m$

In  $\Delta ACB$ ,

$$\tan \alpha = \tan 30^\circ = \frac{AC}{BC} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \frac{BC}{\sqrt{3}} = (AE - CE) = (AE - BD)$$

$$\Rightarrow BC = \sqrt{3} \left( \frac{3 + \sqrt{3}}{2} - 1 \right) h$$

$$\Rightarrow BC = \sqrt{3} \left( \frac{1 + \sqrt{3}}{2} \right) \times 30$$

$$\Rightarrow DE = BC = (45 + 15\sqrt{3})$$

37. (a) Given,  $\beta = 30^\circ$

In  $\Delta ADE$ ,

$$\tan \beta = \frac{AE}{DE}$$

$$\Rightarrow \tan 30^\circ = \left( \frac{\frac{3 + \sqrt{3}}{2}}{DE} \right)$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{\sqrt{3} \left( \frac{1 + \sqrt{3}}{2} \right) h}{DE}$$

$$\Rightarrow DE = \frac{3}{2} (1 + \sqrt{3}) h$$

In  $\Delta BDE$ ,

$$\tan \theta = \frac{BD}{DE} = \frac{h}{DE}$$

$$\Rightarrow \tan \theta = \frac{h}{\frac{3}{2}(1 + \sqrt{3})h}$$

$$\Rightarrow \frac{2}{3} \times \frac{1}{(1 + \sqrt{3})}$$

$$\Rightarrow \frac{2}{3} \times \frac{1}{(1 + \sqrt{3})} \times \frac{(1 - \sqrt{3})}{(1 - \sqrt{3})} = \frac{\sqrt{3} - 1}{3}$$

$$\Rightarrow \frac{(\sqrt{3} - 1)}{3} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{(3 - \sqrt{3})}{3\sqrt{3}}$$

38. (b) Given that,

$$\frac{\tan x}{1 + \sec x} - \frac{\tan x}{1 - \sec x}$$


$$\Rightarrow \frac{\tan x (1 - \sec x - 1 - \sec x)}{1 - \sec^2 x}$$

$$\Rightarrow \frac{-2 \tan x \sec x}{-\tan^2 x} = \frac{2}{\frac{\cos x}{\frac{\sin x}{\cos x}}} = 2 \operatorname{cosec} x$$

39. (b)

$$\frac{\sqrt{\cos x \operatorname{cosec} y - \cos x \sin y}}{\sqrt{\cos x \operatorname{cosec} (90^\circ - x) - \cos x \sin (90^\circ - x)}} \\ \Rightarrow \frac{\sqrt{\cos x \sec x - \cos^2 x}}{\sqrt{1 - \cos^2 x}} = \sqrt{\sin^2 x} = \sin x$$

40. (c)  $\sin \theta = \frac{m^2 - n^2}{m^2 + n^2}$



In  $\Delta ABC$ ,  $BC = \sqrt{(AC)^2 - (AB)^2}$   
 $\frac{\sqrt{m^4 + n^4 + 2m^2n^2 - (m^4 + n^4 - 2m^2n^2)}}{\sqrt{4m^2n^2}} = \frac{2mn}{2mn}$

$$\tan \theta = \frac{m^2 - n^2}{2mn}$$

41. (b)  $1 + \tan \theta = \sqrt{2}$   
 $\Rightarrow \tan \theta = \sqrt{2} - 1$

$$\cot \theta - 1 = \frac{1}{\sqrt{2} - 1} - 1 = \frac{\sqrt{2} + 1}{2 - 1} - 1 \\ \Rightarrow \sqrt{2}$$

42. (c)

$$\sin x : \sin y = \sqrt{3} : 1$$

$$\frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \frac{\sqrt{3}}{1}$$

Hence, we can say  $\sin x = \sin 60^\circ$


$$\sin y = \sin 30^\circ \\ \Rightarrow \frac{x}{y} = \frac{60}{30} = \frac{2}{1}$$

43. (b)

$$\frac{5 \sin 75^\circ \sin 77^\circ + 2 \cos 13^\circ \cos 15^\circ}{\cos 15^\circ \sin 77^\circ} - \frac{7 \sin 81^\circ}{\cos 9^\circ} \\ \Rightarrow \frac{5 \cos 15^\circ \sin 77^\circ + 2 \sin 77^\circ \cos 15^\circ}{\cos 15^\circ \sin 77^\circ} - \frac{7 \cos 9^\circ}{\cos 9^\circ} \\ \Rightarrow \frac{7 \cos 15^\circ \sin 77^\circ}{\cos 15^\circ \sin 77^\circ} - \frac{7 \cos 9^\circ}{\cos 9^\circ} = 7 - 7 = 0$$



44. (c)



$$\cot \theta = \frac{8}{15}$$

$$AC = \sqrt{8^2 + 15^2} = 17$$


$$\Rightarrow \cos \theta = \frac{8}{17}$$

$$\Rightarrow \frac{\sqrt{1 - \cos \theta}}{\sqrt{1 + \cos \theta}} = \frac{\sqrt{1 - \frac{8}{17}}}{\sqrt{1 + \frac{8}{17}}} = \sqrt{\frac{9}{25}} = \frac{3}{5}$$

45. (b) As we know,  $\sin x$  is increasing from 0 to  $90^\circ$

$$\therefore \sin y > \sin x$$

46. (c)



Given,

$$\tan \theta = \frac{p}{q}$$

$$\Rightarrow \sec \theta = \frac{\sqrt{p^2 + q^2}}{q}$$

$$\begin{aligned} \operatorname{cosec} \theta &= \frac{\sqrt{p^2 + q^2}}{p} \\ \frac{p \sec \theta - q \operatorname{cosec} \theta}{p \sec \theta + q \operatorname{cosec} \theta} &= \frac{p \left( \frac{\sqrt{p^2 + q^2}}{q} \right) - q \left( \frac{\sqrt{p^2 + q^2}}{p} \right)}{p \left( \frac{\sqrt{p^2 + q^2}}{q} \right) + q \left( \frac{\sqrt{p^2 + q^2}}{p} \right)} \\ &= \frac{\frac{p}{q} - \frac{q}{p}}{\frac{p}{q} + \frac{q}{p}} = \frac{p^2 - q^2}{p^2 + q^2} \end{aligned}$$


47. (d) We know that, the value of  $\cos \theta$  is decreasing from 0 to  $90^\circ$

$$\therefore \cos 1^\circ > \cos 89^\circ$$

$$\Rightarrow p > q$$

Also,  $\cos 1^\circ$  close to 1 and  $89^\circ$  is close to 0

48. (d)



$$\Rightarrow 7(1 - \sin^2 \theta) + 3(\sin^2 \theta) = 4$$

$$\Rightarrow 7 - 4\sin^2 \theta = 4$$


$$\Rightarrow 4\sin^2 \theta = 3$$

$$\Rightarrow \sin \theta = \pm \frac{\sqrt{3}}{2}$$

$$\text{For } 0 < \theta < \frac{\pi}{2}, \sin \theta = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \tan \theta = \frac{\sqrt{3}}{1} = \sqrt{3}$$

49. (a) Percentage of candidates passed in both the subjects =  $\{100 - (25 + 17 + 35)\}$   
 $\Rightarrow 23\%$



50. (a) Given:  $S = x \in N : \{x + 3 = 3\}$   
 $S = \{\}$ . Thus, S is a null set.

51. (b) Given equation,

$$2x^2 - 3x - 4 = 0$$

For a reciprocal roots, we replace  $x$  by  $1/x$

$$\Rightarrow 2\left(\frac{1}{x}\right)^2 - 3\left(\frac{1}{x}\right) - 4 = 0$$

$$\Rightarrow -4x^2 - 3x + 2 = 0$$

$$\Rightarrow 4x^2 + 3x - 2 = 0$$

52. (c) Given equation is

$$kx^2 + (2k + 6)x + 16 = 0$$

For equal roots then D must be zero

$$\Rightarrow (2k + 6)^2 - 4k \times 16 = 0$$

$$\Rightarrow 4k^2 + 24k + 36 - 64k = 0$$

$$\Rightarrow 4k^2 - 40k + 36 = 0$$

$$\Rightarrow k^2 - 10k + 9 = 0$$

$$\Rightarrow k^2 - 9k - k + 9 = 0$$

$(k - 9)(k - 1)$  Therefore,  $k = 1, k = 9$

53. (c)

Given,  $3^x + 27(3)^{-x} = 12$

Let  $3^x = y$

$$\Rightarrow y + \frac{27}{y} = 12$$

$$\Rightarrow y^2 - 12y + 27 = 0$$

$$\Rightarrow y^2 - 9y - 3y + 27 = 0$$

$$\Rightarrow (y - 3)(y - 9) = 0, y = 3, y = 9$$

54. (c) Let number of student in each row =  $x$   
 and number of rows =  $y$   
 Hence total number of students =  $xy$   
 According to the question,

$$xy = (x + 1)(y - 2)$$

$$\Rightarrow xy = xy - 2x + y - 2$$

$$\Rightarrow 2x - y = -2 \text{ \& } xy = (x - 1)(y + 3)$$

$$\Rightarrow xy = xy + 3x - y - 3$$

$$\Rightarrow -3x + y = 3 \text{ \& } 3x - y = 3$$

Therefore,  $x = 5, y = 12$  \&  $xy = 60$

55. (d) Let Pooja's initial salary is ₹  $x$

Fixed increment every year is ₹  $y$ .

According to question  $x + 3y = 4200$

$$x + 8y = 6800$$

On solving equations (i) and (ii), we get

$$x = ₹ 2640, \quad y = ₹ 520$$

56. (c) Let the two-digit number be  $10y + x$ .

According to question,  $x + y = 10$

$$10y + x - 18 = 10x + y$$

$$\Rightarrow 9x - 9y = -18$$

$$\Rightarrow x - y = -2$$

On solving equations, we get

$$x = 4, \quad y = 6$$

Required product =  $xy = 4 \times 6 = 24$ .

57. (b) Equations  $kx - y = 2, 6x - 2y = 3$  have a unique solution. Then,

$$\frac{k}{6} \neq \frac{1}{2} \Rightarrow k \neq 3$$

58. (c) Given system of equations are:

$$\begin{aligned} x + 2y &= 3 \text{ \& } 3x + 6y = 9 \\ \Rightarrow x + 2y &= 3 \\ \Rightarrow \frac{a_1}{a_2} &= \frac{b_1}{b_2} = \frac{c_1}{c_2} = \frac{1}{3} \end{aligned}$$

So, it has infinitely many solutions

59. (b)

$$\frac{\log_{13}(10)}{\log_{169}(10)} = \frac{\log_{13}(10)}{\log_{13^2}(10)}$$

$$\Rightarrow \frac{\log_{13}10}{\frac{1}{2}\log_{13}10} = \frac{1}{\frac{1}{2}} = 2$$

60. (b)

$$\log_{10} \left[ 1 - \{1 - (1 - x^2)^{-1}\}^{-1} \right]^{-\frac{1}{2}} = 1$$

$$\log_{10} \left[ 1 - \left\{ 1 - \frac{1}{1 - x^2} \right\}^{-1} \right]^{-\frac{1}{2}} = 1$$

$$\log_{10} \left[ 1 - \left\{ \frac{-x^2}{1 - x^2} \right\}^{-1} \right]^{-\frac{1}{2}} = 1$$

$$\log_{10} \left[ 1 - \left\{ \frac{1 - x^2}{-x^2} \right\} \right]^{-\frac{1}{2}} = 1$$

$$\log_{10} \left[ \frac{1}{x^2} \right]^{-\frac{1}{2}} = 1$$

$$\Rightarrow \log_{10} x = 1 \text{ or } \log_{10} x = \log_{10} 10$$

$$\Rightarrow x = 10$$

61. (c)

$$2^x - 2^{x-1} = 4$$

$$\Rightarrow 2^x \left( 1 - \frac{1}{2} \right) = 4$$

$$\begin{aligned} \Rightarrow 2^x &= 8 \Rightarrow 2^x = 2^3 \Rightarrow x = 3 \\ \text{Hence, } 2^x + 2^{x-1} &= 2^3 + 2^2 = 12 \end{aligned}$$

62. (b) Given that,

$$1 + \frac{1}{\left( 1 + \left\{ \frac{1}{\left( 1 + \frac{1}{x} \right)} \right\} \right)} = \frac{11}{7}$$

$$\Rightarrow 1 + \frac{2}{\left\{ 1 + \frac{x}{1+x} \right\}} = \frac{11}{7}$$

$$\Rightarrow 1 + \frac{1+x}{1+2x} = \frac{11}{7}$$

$$\Rightarrow 14 + 21x = 11 + 22x \Rightarrow x = 3$$

63. (b)

$$9\sqrt{2} - \sqrt{8} - 4\sqrt{2}$$

$$\Rightarrow 9\sqrt{2} - 2\sqrt{2} - 4\sqrt{2} = 3\sqrt{2}$$

64. (d)

Given,  $x + \frac{1}{x} = p$

$$\Rightarrow \left( x + \frac{1}{x} \right)^2 = p^2$$

$$\Rightarrow x^2 + \frac{1}{x^2} + 2 = p^2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = p^2 - 2$$

$$\Rightarrow \left( x^2 + \frac{1}{x^2} \right)^3 = (p^2 - 2)^3$$

$$\Rightarrow x^6 + \frac{1}{x^6} + 3 \left( x^2 + \frac{1}{x^2} \right) =$$

$$p^6 - 8 - 6p^2(p^2 - 2)$$

$$\Rightarrow x^6 + \frac{1}{x^6} + 3(p^2 - 2) =$$

$$p^6 - 8 - 6p^4 + 12p^2$$

$$\Rightarrow x^6 + \frac{1}{x^6} = p^6 - 6p^4 + 9p^2 - 2$$

65. (d)

$$\frac{M_1 D_1}{W_1} = \frac{M_2 D_2}{W_2} = \frac{5 \times \frac{1}{2}}{\frac{1}{48}} = \frac{6 \times D_2}{\frac{1}{40}}$$

$$\Rightarrow D_2 = \frac{\frac{1}{2} \times 5 \times 48}{40 \times 6} = \frac{1}{2}$$

66. (a)

$$M_1 D_1 = M_2 D_2$$

$$M_1 = n, D_1 = 20,$$

$$M_2 = (50 + n), D_2 = 16$$

$$\Rightarrow n \times 20 = (n + 50) \times 16$$

$$\Rightarrow 20n = 16n + 800$$

$$\Rightarrow 4n = 800 \Rightarrow n = 200$$

67. (c) According to question.

$$t_1 - t_2 = 12 - (-15) = 27$$

$$\Rightarrow \frac{x}{V_1} - \frac{x}{V_2} = \frac{27}{60}$$

$$\Rightarrow \frac{x}{2.5} - \frac{x}{4} = \frac{27}{60} \Rightarrow \frac{2x}{5} - \frac{x}{4} = \frac{9}{20}$$

$$\Rightarrow \frac{3x}{20} = \frac{9}{20} \Rightarrow 3x = 9 \Rightarrow x = 3 \text{ km}$$

68. (d) Let usual speed =  $v$

Increase speed =  $v + 5$

$$\frac{300}{v} - \frac{300}{v+5} = 2$$

Now,  $\Rightarrow \frac{300(v+5) - 300v}{v(v+5)} = 2$

$$\Rightarrow 2v^2 + 10v - 1500 = 0$$

$$\Rightarrow (v+30)(2v-50) = 0$$

$$\Rightarrow v = 25$$

69. (d) Market price of machine = 18000

Discount of 20% = 3600

Selling Price = 18000 - 3600 = 14400

There was a loss of 4%

$$\Rightarrow x - \frac{4}{100} \times x = 14400$$

$$\Rightarrow \frac{25x - x}{25} = 14400$$

$$\Rightarrow x = \frac{25}{24} \times 14400 = 15000$$

70. (c) Cost price of milk =  $28 \times 8.5 = ₹238$

He added  $x$  ltrs of water

$$(28 + x) \times 8.5 \Rightarrow 238 + 8.5x$$

Profit = SP - CP

$$\Rightarrow 238 + 8.5x - (238) = 8.5x$$

It is said that this  $8.5x$  is 12.5%

$$\Rightarrow \frac{8.5x}{238} \times 100 = 12.5 \Rightarrow 3.5 \text{ ltrs}$$

71. (d) Cost of 2.5 kg rice = ₹125

$$9 \text{ kg rice} = \frac{125}{2.5} \times 9$$

$$\Rightarrow 9 \text{ kg rice} = 4 \text{ kg pulses}$$

$$\Rightarrow 14 \text{ kg pulses} = \frac{125}{2.5} \times \frac{9}{4} \times 14$$

$$\Rightarrow 14 \text{ kg pulses} = 1.5 \text{ kg tea}$$

$$\Rightarrow 2 \text{ kg tea} = \frac{125 \times 9 \times 14 \times 2}{2.5 \times 4 \times 1.5}$$

$$\Rightarrow 2 \text{ kg tea} = 5 \text{ kg nuts}$$

$$\Rightarrow 11 \text{ kg nuts} = \frac{125 \times 9 \times 14 \times 2}{2.5 \times 4 \times 1.5} \times \frac{11}{5}$$

$$\Rightarrow ₹ 4620$$

72. (b) S.I =  $2 \times 10 = 20\%$

$$C.I = x + y + \frac{xy}{100}$$

$$\Rightarrow 10 + 10 + \frac{10 \times 10}{100} = 21\%$$

Difference between interest =  $21 - 20 = 1\%$

$$1\% \times \frac{P}{100} = 10$$

$$P = x = 1000$$

73. (a) Let Principal = P

Interest =  $0.125 \times P$

$$\Rightarrow S.I = \frac{P \times R \times T}{100}$$

$$\Rightarrow 0.125P = \frac{P \times 10 \times T}{100}$$

$$\Rightarrow 1.25 = T$$

$$T = \frac{5}{4} \text{ or } 1\frac{1}{4} \text{ years}$$

74. (c) P = ₹8400 R = 10% n=2

Let instalment = x

$$P = \frac{x}{\left(1 + \frac{R}{100}\right)^1} + \frac{x}{\left(1 + \frac{R}{100}\right)^2}$$

$$\Rightarrow 8400 = \frac{x}{\left(1 + \frac{10}{100}\right)} + \frac{x}{\left(1 + \frac{10}{100}\right)^2}$$

$$\Rightarrow 8400 = \frac{x}{\left(\frac{110}{100}\right)} + \frac{x}{\left(\frac{110}{100}\right)^2}$$

$$\Rightarrow 8400 = x \left[ \frac{10}{11} + \left(\frac{10}{11}\right)^2 \right]$$

$$\Rightarrow 8400 = x \left[ \frac{110 + 100}{121} \right]$$

$$x = \frac{8400 \times 121}{210} = 40 \times 121 = 4840$$

75. (c) Let number of boys and girls in the class be 4x, 3x

Average height of girl is h

$$\Rightarrow 7x \times 4.6 = 4x \times 4.8 + 3x \times h$$

$$\Rightarrow 32.2x = 19.2x + 3x \times h$$

$$\Rightarrow h = \frac{13}{3} = 4.33$$

76. (a) Radha's age = A. Rani's age = B

$$\frac{A - 5}{B - 5} = 3$$

$$\Rightarrow A - 5 = 3(B - 5)$$

$$\frac{A - 1}{B - 1} = 2$$

$$\Rightarrow A - 1 = 2(B - 1)$$

Solving the above two equations we get

$$B = 9, A = 17$$

$$\Rightarrow A - B = 8 \text{ yrs}$$

77. (b) Sum of n consecutive natural numbers

$$= n(n + 1)/2$$

Average of n consecutive natural numbers

$$= (n + 1)/2$$

For first 50 average =  $51/2 = x$

Average of 54 continuous natural number

$$= 55/2$$

$$\Rightarrow \frac{51}{2} + \frac{4}{2} = x + 2$$

78. (a)  $x:y=1:3, y:z=5:k, z:t=2:5$   
 $t:x=3:4$

$$\Rightarrow \frac{x}{y} \times \frac{y}{z} \times \frac{z}{t} \times \frac{t}{x} = 1$$

$$\Rightarrow \frac{1}{3} \times \frac{5}{k} \times \frac{2}{5} \times \frac{3}{4} = 1$$

$$\Rightarrow \frac{1}{2} = k$$

79. (b) Let quantities of milk and water are  $5x, x$  litres

According to the question,

$$\frac{5x}{x+5} = \frac{5}{2}$$

$$\Rightarrow 10x = 5x + 25$$

$$\Rightarrow 5x = 25 \Rightarrow x = 5$$

Quantity of milk in original mixture = 25 l

80. (d) Let initial salary be ₹22x

Final salary be ₹25x

Let initial number of employees = 3y

Final number of employees = 2y

Present bill = Final salary × Final number of employees

$$\Rightarrow 5000 = 25x \times 2y$$

$$\Rightarrow xy = 100$$

Original bill = Initial salary × Initial number of employees

$$22x \times 3y = 66xy$$

$$\Rightarrow 66 \times 100 = ₹6600$$

81. (b)

	X (20%)	Y (10%)
2010	5000	2000

2011	6000	2200
2012	7200	2420
2013	8640	2662

Year in which Difference in price is more than 5000 is 2013

$$\Rightarrow 8640 - 2662 = 5978$$

82. (c) Let initial rent be ₹100 and initial rooms be 100

Initial collection = ₹10000

New rent of 20% increase = 120

New rooms = 120

New collection = 14400

Change in collection = 4400

$$\text{Percentage change} = \frac{4400}{10000} \times 100 = 44\%$$

83. (a) Let income be ₹100

$$\text{Expenditure} = \frac{90}{100} \times 100 = 90$$

Income increased by 20% = 120

New expenditure = 120 - 10 = 110

Change in expenditure = 20

$$\text{Percentage change} = \frac{20}{90} \times 100$$

$$\Rightarrow \frac{200}{9} \text{ or } 22\frac{2}{9}\%$$

84. (c)  $3^x = 4^y = 12^z = k$

$$3 = k^{\frac{1}{x}}, 4 = k^{\frac{1}{y}}, 12 = k^{\frac{1}{z}}$$

$$\Rightarrow 3 \times 4 = 12$$

$$\Rightarrow k^{\frac{1}{x}} + k^{\frac{1}{y}} = k^{\frac{1}{z}}$$

$$\Rightarrow \frac{1}{x} + \frac{1}{y} = \frac{1}{z}$$

$$\Rightarrow \frac{x+y}{xy} = \frac{1}{z} \Rightarrow z = \frac{xy}{x+y}$$

85. (d) Net effect in area =  $200 + 200 + \frac{200 \times 200}{100} = 800\%$

86. (c)  $9^x 3^y = 2187, 2^{3x} 2^{2y} - 4^{xy} = 0$   
 $2^{3x} 2^{2y} = 4^{xy}$

Given,  $3^{2x+y} = 3^7$

$\Rightarrow 2x + y = 7$

Given,  $2^{3x+2y} = 2^{2xy}$

$\Rightarrow 3x + 2y = 2xy$

Solving both equations we get

$\Rightarrow x = 2, y = 3$

Hence,  $x + y = 5$

87. (a)

$= \sqrt[3]{\frac{512}{125}} = \sqrt[3]{\frac{8^3}{5^3}} = \frac{8}{5}$

$\Rightarrow 1 \frac{3}{5}$

88. (a)

$\frac{26-2}{90} - \frac{19-1}{90}$

$\Rightarrow \frac{24}{90} - \frac{18}{90} = \frac{6}{90}$

$\Rightarrow 0.666666... \text{ or } 0.\bar{6}$

89. (c)

$\sqrt{\frac{289}{196}} = 1 + \frac{x}{14} \Rightarrow \frac{17}{14} = 1 + \frac{x}{14}$

$\Rightarrow \frac{x}{14} = \frac{17}{14} - 1 \Rightarrow x = 3$

90. (d)

$\frac{1}{1+\sqrt{2}} + \frac{1}{\sqrt{2}+\sqrt{3}} + \dots + \frac{1}{\sqrt{15}+\sqrt{16}}$

$\frac{1-\sqrt{2}}{1-2} + \frac{\sqrt{2}-\sqrt{3}}{2-3} + \dots + \frac{\sqrt{15}-\sqrt{16}}{15-16}$

$-1(1-\sqrt{2} + \sqrt{2}-\sqrt{3} + \dots + \sqrt{15}-\sqrt{16})$

$\Rightarrow -1(1-\sqrt{16}) = -1(1-4) = 3$

91. (a)

$\frac{(\sqrt{5}-\sqrt{3})^2 - (\sqrt{5}+\sqrt{3})^2}{(\sqrt{5}+\sqrt{3})(\sqrt{5}-\sqrt{3})}$

$\Rightarrow \frac{5+3-2\sqrt{15}-5-3-2\sqrt{15}}{5-3}$

$\Rightarrow \frac{-4\sqrt{15}}{2} = -2\sqrt{15}$

92. (a) Let number be  $x$

According to the question

$x^2 + x = 20$

$\Rightarrow x^2 + x - 20 = 0$

Upon solving  $x = 4, x = -5$

93. (d) We know that LCM is the multiple of HCF. So that 55 cannot be HCF because it is not divisor of 150.

94. (a) For same remainder

$486 - 392 = 94$

$627 - 486 = 141$

$627 - 392 = 235$

HCF of (94, 141, 235) = 47

95. (c) Minimum number of rows

$\Rightarrow \frac{21}{7} + \frac{42}{7} + \frac{56}{7} = 17$

96. (a) Let two digit of the numbers be  $x$  &  $y$

$xy = 8 \Rightarrow y = \frac{8}{x}$

$\Rightarrow (10x + y) = 4(10y + x) + 9$

$$\Rightarrow 10x + y = 40y + 4x + 9$$

$$\Rightarrow 6x = 39y + 9$$

$$\Rightarrow 2x = 13y + 3$$

$$2x = \frac{104}{x} + 3$$

$$\Rightarrow 2x^2 - 3x - 104 = 0$$

$$x = 8, y = 1 \quad \text{Number} = 81$$

97. (c) Quotient = 182,

$$\text{Remainder} = 182 - 175 = 7$$

$$N = 17 \times 182 + 7 = 3101$$

98. (b) Let  $x$  &  $y$  be the number of apples and oranges brought by the person  
 $5x + 7y = 500$

$$y = \frac{500 - 5x}{7} = \frac{5(100 - x)}{7}$$

For  $x$  &  $y$  to be integers

$$x = 2, 9, 16, 23, 30, 37, 46, 51, 58, 65, 72, 79, 86, 93$$

99. (d) Let the three prime numbers be  $x, y, y + 36$

$$x + y + (y + 36) = 100$$

$$x + 2y = 64$$

$2y$  is an even number always as multiplied by 2

We know that even + even = even

Odd + odd = even

So,  $x$  has to be even to satisfy

$$x + 2y = 64$$

The only even prime number is 2

$$x = 2, y = 31$$

Numbers are 2, 31 and 67

100. (c) Dividend =  $Q \times D + R$

$$D = 5Q, D = 2R$$

$$R = 15, D = 30$$

$$5Q = 30, Q = 6$$

$$\text{Dividend} = DQ + R = 30 \times 6 + 15 = 195$$